

# Explosion Energies, Nickel Masses, and Distances of Supernovae of Type IIP

D. K. Nadyozhin

*Institute of Theoretical and Experimental Physics, Moscow, 117259, Russia*

*Max-Planck-Institut für Astrophysik, Garching, 85741, Germany*

*Astronomisches Institut der Universität Basel, Binningen, CH-4102, Switzerland*

Accepted ?????. Received 2003 April 28; in original form 2003 April 28

## ABSTRACT

The hydrodynamical modelling of Type II plateau supernova light curves predicts a correlation between three observable parameters (the plateau duration, the absolute magnitude and photospheric velocity at the middle of the plateau) on the one side and three physical parameters (the explosion energy  $E$ , the mass of the envelope expelled  $\mathcal{M}$ , and the presupernova radius  $R$ ) on the other side. The correlation is used, together with *adopted* EPM distances, to estimate  $E$ ,  $\mathcal{M}$ , and  $R$  for a dozen of well-observed SNe IIP. For this set of supernovae, the resulting value of  $E$  varies within a factor of 6 ( $0.5 \lesssim E/10^{51}\text{erg} \lesssim 3$ ), whereas the envelope mass remains within the limits  $10 \lesssim \mathcal{M}/\mathcal{M}_{\odot} \lesssim 30$ . The presupernova radius is typically  $(200 - 600) R_{\odot}$ , but can reach  $\gtrsim 1000 R_{\odot}$  for the brightest supernovae (e.g., SN 1992am).

A new method of determining the distance of SNe IIP is proposed. It is based on the assumption of a correlation between the explosion energy  $E$  and the  $^{56}\text{Ni}$  mass required to power the post-plateau light curve tail through  $^{56}\text{Co}$  decay. The method is useful for SNe IIP with well-observed bolometric light curves both during the plateau and radioactive tail phases. The resulting distances and future improvements are discussed.


**Key words:** supernovae: general – galaxies: distances and redshifts.

## 1 INTRODUCTION

Plateau Type II supernovae (SNe IIP) are believed to come from the explosion of massive supergiant stars whose envelopes are rich in hydrogen. Their light curves are easy to identify by a long plateau (sometimes up to 120–150 d) which is the result of the propagation of a cooling-and-recombination wave (CRW) through the supernova envelope that is in a state of free inertial expansion ( $u = r/t$ ). The CRW physics is discussed in detail by Imshennik & Nadyozhin (1964), Grassberg, Imshennik & Nadyozhin (1971), and Grassberg & Nadyozhin (1976). The CRW propagates supersonically downward through the expanding supernova envelope and separates almost recombined outer layers from still strongly ionized inner ones. During the plateau phase, the photosphere sits on the upper edge of the CRW front. Since the CRW downward speed turns out to be close to the velocity of the outward expansion, the photospheric radius changes only slowly during the plateau phase. If one takes into account that also the effective temperature does not change appreciably (it approximately equals the recombination temperature 5000–7000 K), the approximate constancy of the luminosity becomes obvious.

The supernova outburst properties are determined mainly by three physical parameters: the explosion energy  $E$ , the mass  $\mathcal{M}$  of the envelope expelled, and the initial radius  $R$  of the star just before the explosion (presupernova). Litvinova & Nadyozhin (1983, 1985) have undertaken an attempt to derive these parameters from a comparison of the hydrodynamical supernova models with observations. They constructed simple approximation formulae which allow to estimate  $E$ ,  $\mathcal{M}$ , and  $R$  from the observations of individual SNe IIP. Their results were confirmed by an independent semi-analytical study (Popov 1993). At that time, only one or two supernovae were sufficiently observed to apply these formulae. At present, there exist detailed observational data for 14 such supernovae, including in 12 cases expanding photosphere (EPM) distances, which we use in section 2 to estimate  $E$ ,  $\mathcal{M}$ , and  $R$  by means of these formulae.

In section 3, we propose a new method of distance determination and employ it to 9 individual SNe IIP which are well-observed both at the plateau and radioactive-tail phases. The method is based on the assumption of a correlation between the explosion energy  $E$  and the mass of  $^{56}\text{Ni}$  in the supernova envelope. In section 4 we compare physical parameters and distances of SNe IIP as derived from



**Figure 1.** A schematic SN IIP light curve. An open circle marks the middle of the plateau and two black circles show the plateau boundaries. The light curve tail powered by the  $^{56}\text{Co}$  decay is also shown ( $\tau_{\text{Co}} = 111.3$  d).

the new method with those obtained previously from the EPM method and discuss also other aspects of our results. Concluding remarks are given in section 5.

The preliminary results of this study were reported to the Workshop on physics of supernovae held at Garching, Germany, July 2002 (Nadyozhin 2003).

## 2 A COMPARISON OF HYDRODYNAMIC MODELS WITH OBSERVATIONS

Figure 1 shows a schematic SNe IIP light curve. The plateau is defined as part of the light curve on which the supernova brightness remains within 1 mag of the mean value. For some supernovae, the plateau begins almost immediately after the onset of the explosion ( $t = 0$ ) whereas for others a short luminosity peak can precede the plateau. The peak either appears owing to a shock wave breakout in the case of presupernovae of not very large initial radii ( $R \lesssim 1000 R_{\odot}$ ) or, according to Grassberg et al. (1971), originates from the emergence of a thermal wave precursor for presupernovae of very large radii ( $R \approx (2000 - 5000) R_{\odot}$ ) and of moderate explosion energies ( $E \lesssim 1 \times 10^{51}$  erg), or at last it may occur as a result of interaction between the supernova envelope and a dense stellar wind (Grassberg & Nadyozhin 1987). For some SNe IIP the peak duration  $\delta t$  lasts only a few days and is difficult to observe (shock wave breakout), for others it could be as large as 10–20 days (thermal wave or dense wind) – examples for the latter may be such supernovae as SNe 1988A, 1991al, and 1992af (see below).

It is quite clear that the middle of the plateau is to be used as the main reference point to compare the theoretical models with observations. Litvinova & Nadyozhin (1983, 1985) (LN83 and LN85, hereafter) calculated a grid of the supernova models for  $E$ ,  $\mathcal{M}$ , and  $R$  within limits of  $(0.18 - 2.91) \times 10^{51}$  erg,  $(1 - 16) \mathcal{M}_{\odot}$ , and  $(300 - 5000) R_{\odot}$ . They found  $E$ ,  $\mathcal{M}$ ,  $R$  to be strongly correlated with the plateau duration  $\Delta t$ , and the middle-plateau values of the absolute  $V$ -magnitude  $M_V$ , and the expansion velocity  $u_{\text{ph}}$  at the level of the photosphere (Fig. 1). According to LN85, the following approximate relations can be used to derive  $E$ ,  $\mathcal{M}$ , and  $R$  from observations:

$$\lg E = 0.135 M_V + 2.34 \lg \Delta t + 3.13 \lg u_{\text{ph}} - 4.205, \quad (1)$$

$$\lg \mathcal{M} = 0.234 M_V + 2.91 \lg \Delta t + 1.96 \lg u_{\text{ph}} - 1.829, \quad (2)$$

$$\lg R = -0.572 M_V - 1.07 \lg \Delta t - 2.74 \lg u_{\text{ph}} - 3.350, \quad (3)$$

where  $E$  is expressed in units of  $10^{51}$  erg,  $\mathcal{M}$  and  $R$  are in solar units,  $\Delta t$  in days, and  $u_{\text{ph}}$  in  $1000 \text{ km s}^{-1}$ . Here  $M_V$  can be expressed through the apparent  $V$  magnitude by the relation:

$$M_V = V - A_V - 5 \lg(D/1\text{Mpc}) - 25, \quad (4)$$

where  $D$  is the distance to a supernova and  $A_V$  is the total absorption on the way to the supernova. One can find from Eqs. (1)–(3) that  $E$ ,  $\mathcal{M}$ , and  $R$  scale with the distance as:

$$E \sim D^{-0.675}, \quad \mathcal{M} \sim D^{-1.17}, \quad R \sim D^{2.86}. \quad (5)$$

Thus, it is very important to know  $D$  with as high accuracy as possible. We have selected 14 SNe whose observational data are collected in Table 1. The entries are: the heliocentric recession velocities  $v_0$  (from the NED: NASA/IPAC Extragalactic Database) in column 3, the total absorption  $A_V$  in column 4, the apparent  $V$  magnitude of the mid-point of the plateau in column 5, the duration  $\Delta t$  of the plateau in column 6, and the photosphere expansion velocity  $u_{\text{ph}}$  in column 7. The references are in column 8.

In order to check the extrapolative capability of Eqs. (1)–(3), we have included the SN 1987A in our analysis. It is well known, that the SN 1987A presupernova radius was as small as  $\approx 50 R_{\odot}$  – i.e. outside the interval of  $(300 - 5000) R_{\odot}$  encompassed by the above equations. Moreover, the major part of the SN 1987A plateau (about 70 of 110 d) was powered by the  $^{56}\text{Co}$ -decay (see the review of Imshennik & Nadyozhin 1989 and references therein).

Derived properties of the 14 SNe IIP are in Table 2. Column 2 is the recession velocity  $v_{220}$  of the supernova (column 1) corrected for a self-consistent Virgocentric infall model with a local infall vector of  $220 \text{ km s}^{-1}$  as described by Kraan-Korteweg (1986). Column 3 gives the distance  $D_H = v_{220}/H_0$  assuming arbitrarily a value of  $H_0 = 60 \text{ km s}^{-1} \text{ Mpc}^{-1}$ . For comparison, column 4 gives the distance  $D_{\text{EPM}}$  obtained with the use of the expanding photosphere method in the references listed at the bottom of the table. The SNe 1991al and 1992af are the exception. Owing to the incompleteness of the observational data, it is hard to determine the EPM distance to the SN 1991al (Hamuy 2001). For the same reason, the EPM distance of 55 Mpc for the SN 1992af obtained by Schmidt et al. (1994a) seems to be quite uncertain as pointed out by Hamuy (2001). For these two SNe, we present in column 4 the distances calculated by Hamuy (2001) from the CMB redshifts and the Hubble constant  $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ . Columns 5–8 are the absolute magnitude  $M_V$  of the mid-point of the plateau, the explosion energy  $E$ , the mass expelled  $\mathcal{M}$ , and the presupernova radius  $R$  – all derived from Eqs. (4), (1)–(3) for the  $D_H$  distances listed in column 3. Column 9 gives the mass of  $^{56}\text{Ni}$ , ejected by some supernovae, which was estimated by reducing the radioactive-tail luminosities, measured by Hamuy (2001), to the distances  $D_H$  given in column 3.

For SN 1987A, the resulting values of  $E$  and  $\mathcal{M}$  (Table 2) differ no more than by a factor of 1.5 from current estimates based on a detailed study. However, the presupernova radius turned out to be too large. This happened because the LN85 approximations do not take into account the radioactive heating. An advanced study (Grassberg &

**Table 1.** Observational data for 14 SNe IIP.

| SN     | Host galaxy  | $v_0$<br>km s <sup>-1</sup> | $A_V$<br>mag | $V$<br>mag | $\Delta t$<br>days | $u_{\text{ph}}$<br>km s <sup>-1</sup> | Ref.              |
|--------|--------------|-----------------------------|--------------|------------|--------------------|---------------------------------------|-------------------|
| (1)    | (2)          | (3)                         | (4)          | (5)        | (6)                | (7)                                   | (8)               |
| 1968L  | NGC 5236     | 516                         | 0.219        | 12.0       | 80                 | 4100                                  | 1, 2, 3           |
| 1969L  | NGC 1058     | 518                         | 0.203        | 13.4       | 100                | 4000                                  | 1, 2              |
| 1986L  | NGC 1559     | 1292                        | 0.099        | 14.7       | 110                | 4000                                  | 4                 |
| 1988A  | NGC 4579     | 1519                        | 0.136        | 15.0       | 110                | 3000                                  | 1, 2, 4, 5, 6     |
| 1989L  | NGC 7339     | 1313                        | 1.00         | 16.5       | 140                | 3000                                  | 7, 19             |
| 1990E  | NGC 1035     | 1241                        | 1.083        | 16.0       | 120                | 4000                                  | 2, 4, 8, 9        |
| 1991al | LEDA 140858  | 4572                        | 0.318        | 17.0       | 90                 | 6000                                  | 4                 |
| 1992af | ESO 340-G038 | 6000                        | 0.171        | 17.3       | 90                 | 6000                                  | 4, 7              |
| 1992am | anon 0122-04 | 14600                       | 0.464        | 19.0       | 110                | 4800                                  | 4, 10             |
| 1992ba | NGC 2082     | 1104                        | 0.193        | 15.43      | 100                | 2900                                  | 4, 7              |
| 1999cr | ESO 576-G034 | 6069                        | 0.324        | 18.6       | 100                | 3600                                  | 4                 |
| 1999em | NGC 1637     | 717                         | 0.314        | 14.0       | 110                | 3000                                  | 4, 11, 12, 13, 17 |
| 1999gi | NGC 3184     | 592                         | 0.65         | 15.0       | 110                | 2900                                  | 14, 15, 16, 18    |
| 1987A  | LMC          | 278                         | 0.465        | 3.3        | 110                | 2900                                  | 4                 |

REFERENCES. – (1) Patat et al. 1993; (2) Schmidt et al. 1992; (3) Wood & Andrews 1974; (4) Hamuy 2001; (5) Ruiz-Lapuente et al. 1990; (6) Turatto et al. 1993; (7) Schmidt et al. 1994a; (8) Schmidt et al. 1993; (9) Benetti et al. 1994; (10) Schmidt et al. 1994b; (11) Hamuy et al. 2001; (12) Haynes et al. 1998; (13) Baron et al. 2000; (14) Schlegel 2001; (15) Smartt et al. 2001; (16) Li et al. 2002; (17) Elmhamdi et al. 2003; (18) Leonard et al. 2002b; (19) Pennypacker & Perlmutter 1989

**Table 2.** The supernova physical properties.

| SN     | $v_{220}$<br>km s <sup>-1</sup> | $D_H$<br>Mpc | $D_{\text{EPM}}$<br>Mpc | $M_V$<br>mag | $E$<br>10 <sup>51</sup> erg | $\mathcal{M}$<br>$\mathcal{M}_{\odot}$ | $R$<br>$R_{\odot}$ | $\mathcal{M}_{\text{Ni0}}$<br>$\mathcal{M}_{\odot}$ |
|--------|---------------------------------|--------------|-------------------------|--------------|-----------------------------|----------------------------------------|--------------------|-----------------------------------------------------|
| (1)    | (2)                             | (3)          | (4)                     | (5)          | (6)                         | (7)                                    | (8)                | (9)                                                 |
| 1968L  | 291                             | 4.85         | 4.5 <sup>(1)</sup>      | -16.65       | 0.83                        | 10.3                                   | 286                |                                                     |
| 1969L  | 766                             | 12.77        | 10.6 <sup>(1)</sup>     | -17.33       | 1.05                        | 13.0                                   | 595                |                                                     |
| 1986L  | 1121                            | 18.68        | 16.0 <sup>(1)</sup>     | -16.76       | 1.56                        | 23.5                                   | 251                | 0.026                                               |
| 1988A  | 1179                            | 19.65        | 20.0 <sup>(1)</sup>     | -16.60       | 0.67                        | 14.5                                   | 452                | 0.082                                               |
| 1989L  | 1556                            | 25.93        | 17.0 <sup>(1)</sup>     | -16.57       | 1.18                        | 29.8                                   | 334                |                                                     |
| 1990E  | 1238                            | 20.63        | 18.0 <sup>(1)</sup>     | -16.66       | 1.98                        | 31.9                                   | 200                | 0.052                                               |
| 1991al | 4476                            | 74.60        | 70.0 <sup>(2a)</sup>    | -17.68       | 2.61                        | 17.6                                   | 347                | 0.12                                                |
| 1992af | 6000                            | 100.00       | 83.70 <sup>(2a)</sup>   | -17.87       | 2.46                        | 15.9                                   | 445                | 0.24                                                |
| 1992am | 14600                           | 243.33       | 180.0 <sup>(1)</sup>    | -18.40       | 1.66                        | 13.9                                   | 1321               | 0.36                                                |
| 1992ba | 1096                            | 18.27        | 22.0 <sup>(2)</sup>     | -16.07       | 0.57                        | 13.7                                   | 272                | 0.029                                               |
| 1999cr | 6069                            | 101.15       | 86.0 <sup>(2)</sup>     | -16.75       | 0.90                        | 14.5                                   | 368                | 0.085                                               |
| 1999em | 743                             | 12.38        | 8.20 <sup>(3)</sup>     | -16.78       | 0.63                        | 13.2                                   | 569                | 0.058                                               |
| 1999gi | 707                             | 11.78        | 11.10 <sup>(4)</sup>    | -16.01       | 0.72                        | 18.7                                   | 226                | 0.025*                                              |
| 1987A  | —                               | 0.05         | 0.05                    | -15.66       | 0.80                        | 22.6                                   | 143                | 0.065                                               |

<sup>(1)</sup>Eastman, Schmidt, & Kirshner 1996; <sup>(2)</sup>Hamuy 2001; <sup>(2a)</sup>Hamuy 2001, based on an adopted value of  $H_0 = 65$ , see text; <sup>(3)</sup>Leonard et al. 2002a;

<sup>(4)</sup>Leonard et al. 2002b

\*Derived from  $V = 17.86$  mag at  $t = 174.3$  d (Leonard et al. 2002b), and Hamuy’s recipe (section 5.3 of his Thesis) to convert  $V$  into luminosity  $L$ .

Nadyozhin 1986) demonstrates that the radioactive heating influences only weakly on  $E$  and  $\mathcal{M}$  furnished by Eqs. (1)–(3), whereas the  $R$ -values can be overestimated by a factor of 3. In this connection, one should have in mind that for some supernovae the  $R$ -values from Table 2 can be larger than actual presupernova radii.

According to Table 2, the resulting values of  $E$ ,  $\mathcal{M}$ , and  $R$  seem to be reasonable enough: the expelled mass,

explosion energy, and presupernova radius remain approximately in limits  $(10 - 30) \mathcal{M}_{\odot}$ ,  $(0.6 - 2.6) \times 10^{51}$  erg, and  $(200 - 1300) R_{\odot}$ , respectively. Hamuy (2001) assumed that SNe 1991al, 1992af were discovered several weeks after the explosion. Their plateaus, therefore, could have lasted for  $\Delta t \approx 110$  d. It is quite probable, however, that their peak duration was  $\delta t \approx 20$  d for the reasons mentioned above. Having this in mind, we have chosen in Table 1  $\Delta t = 90$  d

which results in  $u_{\text{ph}} = 6000 \text{ km s}^{-1}$ . In the case of  $\Delta t = 110 \text{ d}$  we would have to assume  $u_{\text{ph}} = 7000 \text{ km s}^{-1}$  and would obtain very large values of  $E$  and  $\mathcal{M}$  for both supernovae:  $E \approx 7 \times 10^{51} \text{ erg}$  and  $\mathcal{M} \approx 40 \mathcal{M}_{\odot}$ . No other special adjustments of the observational data given in Table 1 were made.

### 3 PLATEAU-TAIL DISTANCE DETERMINATION

The SN IIP light curve tails are believed to be powered by the  $^{56}\text{Co}$  decay. The temporal behavior of the bolometric luminosity is given by (see, e.g. Nadyozhin 1994):

$$L = 1.45 \times 10^{43} \exp\left(-\frac{t}{\tau_{\text{Co}}}\right) \frac{\mathcal{M}_{\text{Ni0}}}{\mathcal{M}_{\odot}} \text{ erg s}^{-1}, \quad (6)$$

where  $t$  is measured from the moment of explosion ( $t = 0$ ),  $\mathcal{M}_{\text{Ni0}}$  is the total mass of  $^{56}\text{Ni}$  at  $t = 0$  which decays with a half-life of 6.10 d into  $^{56}\text{Co}$ , and  $\tau_{\text{Co}} = 111.3 \text{ d}$ .

Equation (6) can be written in the form

$$\mathcal{M}_{\text{Ni0}} = \frac{D^2}{145} Q, \quad Q \equiv F_{41}(t) \exp\left(\frac{t}{111.3}\right), \quad (7)$$

where  $\mathcal{M}_{\text{Ni0}}$  is in  $\mathcal{M}_{\odot}$ ,  $t$  in days and  $D$  in Mpc. The quantity  $F_{41}(t)$  is the bolometric tail luminosity measured at time  $t$  in units  $10^{41} \text{ erg s}^{-1}$  under the assumption that the supernova is at distance  $D = 1 \text{ Mpc}$ . Equation (7) contains a single observational parameter  $Q$  which is independent of  $t$  and also makes no assumption on  $D$  as long as  $F_{41}$  is fixed by observations.

Thus, it is irrelevant at which  $t$  the luminosity is actually measured – one has only to be sure that the supernova really entered its tail phase. Columns 8 – 10 of Table 3 give  $t$  and corresponding values of  $F_{41}(t)$  and  $Q$  derived from Hamuy's (2001) Figures 5.7 and 5.8 except SN 1999gi for which the values were calculated from the data of Leonard et al. (2002b).

If the value of  $\mathcal{M}_{\text{Ni0}}$  was known, one could easily find the distance  $D$  from Eq. (7). So, we have to look for a way to estimate  $\mathcal{M}_{\text{Ni0}}$  independently. It seems reasonable to assume that the supernova explosion energy  $E$  should correlate with  $\mathcal{M}_{\text{Ni0}}$  produced during the explosion. This means that

$$E = f(\mathcal{M}_{\text{Ni0}}) = f\left(\frac{D^2}{145} Q\right), \quad (8)$$

where  $f$  represents a *statistically admissible correlation function* rather than a strict mathematical relation. Inserting this expression for  $E$  into Eq. (1) and using Eq. (4) for  $M_V$ , we obtain an equation which can be solved for  $D$  when  $V - A_V$ ,  $u_{\text{ph}}$ ,  $\Delta t$ , and  $Q$  are known from observations. Then for given  $D$ , we can find  $E$ ,  $\mathcal{M}$ ,  $R$ , and  $\mathcal{M}_{\text{Ni0}}$  from Eqs. (1)–(3), Eq. (4), and Eq. (7), respectively.

What can be said about the function  $f(\mathcal{M}_{\text{Ni0}})$  at present, when the details of the SN II mechanism remain still ambiguous? First of all, it is reasonable to assume that a good fraction of  $E$  comes from the recombination of free neutrons and protons into  $^{56}\text{Ni}$  just at the bottom of the envelope to be finally expelled (Nadyozhin 1978, Bethe 1996). The hydrodynamical modelling of the collapse (Nadyozhin 1978) have indicated that under favourable conditions a neutron-proton shell could be accumulated just under the

steady accreting shock wave. When the mass of such a shell reaches some critical value (presumably of the order of  $\approx 0.1 \mathcal{M}_{\odot}$ ) the shell can become unstable in respect to recombining into the "iron group" elements (specifically into  $^{56}\text{Ni}$ ) to supply the stalled shock wave with the energy of  $\approx 10^{51} \text{ erg}$  necessary to trigger the supernova. Here, there is a physical analogy with the origin of planetary nebulae from red giants where the energy from the recombination of hydrogen and helium causes the expulsion of a red giant rarefied envelope. The recent study (Imshennik 2002, and references therein) of the "neutrino crown" – the region enclosed within neutrinosphere and accreting shock, turns out to be in line with such a picture of the supernova mechanism. However, some Ni can be produced through the explosive carbon-oxygen burning induced by the outgoing shock wave. In this case the energy release per unit Ni mass is lower by an order of magnitude than for the neutron-proton recombination.

The energy released by the neutron-proton recombination, producing a  $^{56}\text{Ni}$  mass of  $\mathcal{M}_{\text{Ni0}}$ , is given by

$$E(\text{np} \rightarrow \text{Ni}) = 1.66 \times 10^{52} \frac{\mathcal{M}_{\text{Ni0}}}{\mathcal{M}_{\odot}} \text{ erg}. \quad (9)$$

Thus, the production of only  $\sim 0.06 \mathcal{M}_{\odot}$  of  $^{56}\text{Ni}$  is sufficient to provide the standard explosion energy of  $10^{51} \text{ erg}$ . The current hydrodynamic models of the SN II explosions (Woosley & Weaver 1995; Rauscher et al. 2002) do not show a correlation between  $E$  and  $\mathcal{M}_{\text{Ni0}}$  because in these models  $^{56}\text{Ni}$  comes from explosive silicon and carbon-oxygen burning near to the envelope bottom and its yield is sensitive to the mass cut point. The photometrical and spectroscopical properties of the SN models are virtually independent of the mass cut. On the contrary, the nucleosynthesis yields are very sensitive to the mass cut. In the current SN models the explosion is usually simulated by locating a piston at the internal boundary  $m = \mathcal{M}_{\text{cut}}$ . The piston moves with time according to a prescribed law,  $R_{\text{pis}}(t)$ , with the velocity ( $\dot{R}$ ) amplitude being chosen to ensure the final kinetic energy of the expelled envelope of the order of  $10^{51} \text{ erg}$ . There are two major uncertainties at this point. First, for a given velocity amplitude the resulting nuclear yields are still sensitive to the form of the function  $R_{\text{pis}}(t)$ . Second, the presupernova structure (especially chemical composition) in the vicinity of  $m = \mathcal{M}_{\text{cut}}$  will always remain ambiguous until the detailed mechanism of the SN disintegration onto the collapsed core and thrown envelope is established. The point is that such 2D effects as rotation and large-scale mixing can result in the presupernova structure different from that predicted by the spherically symmetrical models. Under such circumstances, it is difficult to find a serious argument against the possibility to expel a noticeable amount of  $^{56}\text{Ni}$  from the recombination of the neutron-proton shell. Thus, we propose a neutron-proton layer which is located somewhat deeper than the value of  $\mathcal{M}_{\text{cut}}$  assumed in the current SN models. This layer recombines into  $^{56}\text{Ni}$  providing the energy sufficient to convert a steady-state accretion shock into the outgoing blast wave. In this case a good correlation between  $E$  and  $\mathcal{M}_{\text{Ni0}}$  is to be expected.

The proposed correlation can have a complex nature. It is quite probable that the function  $f$  in Eq. (8) depends also on  $\mathcal{M}$  since the supernova mechanism is expected to be sensitive to the presupernova mass. For us only the existence

**Table 3.** The tail-calibrated supernova physical properties ( $\xi = 1$ ).

| SN     | $D_{P-T}$<br>Mpc | $M_V$<br>mag | $E$<br>$10^{51}$ erg | $\mathcal{M}$<br>$\mathcal{M}_{\odot}$ | $R$<br>$R_{\odot}$ | $\mathcal{M}_{Ni0}$<br>$\mathcal{M}_{\odot}$ | $F_{41}(t)$<br>$10^{41}$ erg/(s Mpc <sup>2</sup> ) | $t$<br>days | $Q$     |
|--------|------------------|--------------|----------------------|----------------------------------------|--------------------|----------------------------------------------|----------------------------------------------------|-------------|---------|
| (1)    | (2)              | (3)          | (4)                  | (5)                                    | (6)                | (7)                                          | (8)                                                | (9)         | (10)    |
| 1986L  | 29.67            | −17.76       | 1.14                 | 13.7                                   | 944                | 0.067                                        | $2.25 \times 10^{-3}$                              | 180         | 0.0113  |
| 1988A  | 15.21            | −16.05       | 0.79                 | 19.6                                   | 217                | 0.048                                        | $4.96 \times 10^{-3}$                              | 200         | 0.0299  |
| 1990E  | 29.16            | −17.41       | 1.57                 | 21.3                                   | 539                | 0.094                                        | $2.67 \times 10^{-3}$                              | 200         | 0.0161  |
| 1991al | 85.31            | −17.97       | 2.38                 | 15.0                                   | 509                | 0.14                                         | $8.13 \times 10^{-4}$                              | 140         | 0.00286 |
| 1992af | 86.45            | −17.55       | 2.71                 | 18.8                                   | 293                | 0.16                                         | $9.02 \times 10^{-4}$                              | 140         | 0.00317 |
| 1992ba | 19.85            | −16.25       | 0.53                 | 12.4                                   | 346                | 0.032                                        | $1.97 \times 10^{-3}$                              | 200         | 0.0119  |
| 1999em | 11.08            | −16.54       | 0.68                 | 15.0                                   | 414                | 0.041                                        | $1.26 \times 10^{-2}$                              | 150         | 0.0485  |
| 1999gi | 14.53            | −16.46       | 0.63                 | 14.5                                   | 411                | 0.038                                        | $5.41 \times 10^{-3}$                              | 174         | 0.0259  |
| 1987A  | 0.045            | −15.42       | 0.87                 | 25.6                                   | 104                | 0.053                                        | $8.16 \times 10^2$                                 | 170         | 3762    |

of some correlation is important which in combination with Eqs. (1)–(3) allows to determine the distance independently.

To demonstrate how such a method can work we make the simplest assumption that  $E$  is proportional to  $E(\text{np} \rightarrow \text{Ni})$ . Then one can write:

$$E = \xi E(\text{np} \rightarrow \text{Ni}) = 16.6 \xi \mathcal{M}_{Ni0} = 0.1145 \xi D^2 Q, \quad (10)$$

where, as usual,  $E$  is in  $10^{51}$  erg,  $\mathcal{M}_{Ni0}$  in  $\mathcal{M}_{\odot}$  and  $D$  in Mpc. This equation implies that the function  $f$ , introduced in Eq. (8), reads as  $f(x) = 16.6 \xi x$  where  $\xi$  is an adjustable parameter which can be either less or larger than 1. If there is a noticeable contribution to  $\mathcal{M}_{Ni0}$  from the explosive carbon-oxygen burning then  $\xi < 1$ ; if a noticeable contribution to the explosion energy comes from other source rather than the neutron-proton recombination then  $\xi > 1$ .

Inserting  $E$  from Eq. (10) and  $M_V$  from Eq. (4) into Eq. (1) and solving for  $D$ , we obtain:

$$\lg D = -0.374 \lg(\xi Q) + 0.0504 (V - A_V) + 0.875 \lg \Delta t + 1.17 \lg u_{ph} - 2.482, \quad (11)$$

where  $D$  is in Mpc,  $\Delta t$  in days, and  $u_{ph}$  in  $1000 \text{ km s}^{-1}$ . We will refer to distances derived from Eq. (11) as ‘plateau-tail distances’,  $D_{P-T}$ , hereafter. The results are given in Table 3 for nine supernovae selected from Table 2. We did not include SNe 1992am and 1999cr in our analysis because their last available observations may not yet reflect the radioactive tail phase. Specifically, there are only two observations of SN 1992am at the post-plateau phase of the light curve. Since the observations are separated by a short time interval of 3 days, it is difficult to derive the inclination of the bolometric light curve with a required accuracy to be sure that SN 1992am is already in the radioactive-tail phase. Moreover, one has to remember that in addition to the Co-decay the tail luminosity can also be contributed by the ejecta-wind interaction (see Chugai 1991 and references therein). SN 1992am is suspicious in this respect because its presupernova radius seems to be larger than  $1000 R_{\odot}$  (Table 2). Hence, the  $\mathcal{M}_{Ni0}$ -values for these SNe in Table 2 could be actually upper limits.

The different columns of Table 3 give the following quantities: (2) the distance  $D_{P-T}$  from Eq. (11) setting  $\xi = 1$ ; (3) the corresponding absolute  $V$ -magnitude of the mid-point of the plateau  $M_V$ ; (4)–(7) the quantities  $E$ ,  $\mathcal{M}$ ,  $R$ , and  $\mathcal{M}_{Ni0}$  as in Table 2, but now using the distance  $D_{P-T}$  as in column (1); the columns (8)–(10) are explained above.

The values of  $E$ ,  $\mathcal{M}$ ,  $R$ , and  $\mathcal{M}_{Ni0}$  for the  $\xi$ -values different from 1 can be found using the following scaling relations which result from Eqs. (5), (7), and (11):


$$E \sim \xi^{0.252}, \mathcal{M} \sim \xi^{0.438}, R \sim \xi^{-1.07}, \mathcal{M}_{Ni0} \sim \xi^{-0.748}. \quad (12)$$

For a fixed  $Q$ , the dependence of the distance  $D_{P-T}$ , defined by Eq. (11), on extinction  $A_V$  proves to be very weak: an error in  $A_V$  of  $\pm 1$  mag changes  $D_{P-T}$  by only  $\pm 12\%$ . However, if the tail luminosity  $F_{41}$  is derived from the  $V$  measurements (just the case of Hamuy’s  $F_{41}$ -values we use here) then the  $\lg F_{41}$ , and consequently  $\lg Q$ , scale as  $0.4 A_V$  and  $\lg D_{P-T}$ , derived from Eq. (11), actually varies with  $A_V$  in a standard way, as  $-0.2 A_V$ . If the tail luminosity were derived from infra-red measurements then the resulting  $D_{P-T}$  distances would be largely independent of extinction. Note also rather weak dependence on  $\xi Q$ :  $D_{P-T} \sim (\xi Q)^{-0.374}$ . For instance, the decrease in  $\xi Q$  by a factor of 2 results in an *increase* of  $D_{P-T}$  by 30% only.

The random errors typically of  $\pm 10\%$  for the  $\delta t$  and  $u_{ph}$  values assumed in Table 1 result in the uncertainty factor of  $\approx 1.2$  for  $D_{P-T}$  and  $\approx 1.5$  for  $\mathcal{M}_{Ni0}(\sim D^2)$  given in Table 3. However, one has to keep in mind two main sources of systematic errors: (i) probable deviation of the theoretical models (which Eqs. 1–3 are based on) from real SNe, and (ii) the presentation of the  $E - \mathcal{M}_{Ni0}$  correlation in the form of the straight proportionality (Eq. 10). Both the types of systematic errors are difficult to estimate at present. Although the SN models calculated in LN83 and LN85 rest upon a very simplified presupernova structure, they consistently take into account the ionization and recombination of hydrogen and helium thereby remaining still useful. When a new grid of the SN models, based on modern evolutionary presupernova structure, is created the systematic error (i) certainly will be reduced. The reduction of the systematic error (ii) requires a more profound knowledge of the SN mechanism. Empirically, this problem can be solved by adjusting the factor  $\xi$  for each individual SN. It is necessary, however, to collect a much more rich statistics (at least by a factor of 3) than that available nowadays (only 9 SNe in Table 3).

## 4 DISCUSSION

The plateau-tail distances derived in section 3 and listed in column 2 of Table 3 are plotted in a Hubble diagram in Fig. 2




**Figure 2.** The Hubble diagram of 8 SNe IIP with  $D_{P-T}$  distances from plateau and tail observations (black circles). Also shown are the eleven SNe IIP with known EPM distances (asterisks). The respective Hubble lines are fitted to the data. The abscissa is the distance modulus  $(m - M) = 5 \lg D + 25$ .


(except SN 1987A which is not in the Hubble flow). The eight SNe IIP define a Hubble line with  $H_0 = 55 \pm 5 \text{ km s}^{-1} \text{ Mpc}^{-1}$ . Also shown in Fig. 2 are the eleven SNe IIP for which EPM distances have been published (column 3 of Table 2). They define a Hubble line of  $H_0 = 70 \pm 4 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , i.e. the EPM distances are smaller than the plateau-tail distances by 25% on average.

At this point it is not possible to decide which of the two results is more nearly correct. Both methods, the plateau-tail distances and the EPM distances, depend on assumptions which are difficult to verify. The EPM method faces the problem of the dilution factor in an expanding atmosphere and the definition of the photospheric radius which depends on the uncertainties connected with the opacity of an expanding medium. However, it may be noted that the EPM distance of SN 1987A agrees well with the generally adopted distance of LMC of 50 kpc (Eastman, Schmidt, & Kirshner 1996) and the EPM distance of SN 1968L is indistinguishable from the Cepheid distance of NGC 5236 (M83) (Thim et al. 2003).

The main assumption which affects the plateau-tail distances concerns the nature of the proposed  $E - \mathcal{M}_{\text{NiO}}$  correlation. For our simplified example of such a correlation, all the uncertainties turn out to be cumulated in the proportionality factor  $\xi$  between the explosion energy  $E$  and the nickel mass  $\mathcal{M}_{\text{NiO}}$ . In Table 3 we have adopted a plausible value of  $\xi = 1$ , but it cannot be excluded that  $\xi$  is as low as 0.5 or as high as 2. Since the Hubble constant scales as  $H_0 \sim \xi^{0.374}$ , an average value as high as  $\xi = 1.9$  would be needed to bring the plateau-tail distances in general accord with the EPM distances. Such a high average value of  $\xi$  is, however, not supported by SNe 1987A and 1999gi. If the  $D_{P-T}$  distance of SN 1987A from Table 3 is scaled to the canonical LMC distance of 50 kpc,  $\xi$  becomes 0.75. And if the host galaxy NGC 3184 of SN 1999gi with a  $D_{P-T}$



**Figure 3.** The explosion energies  $E(H_0 = 60)$  and  $E(\text{EPM})$  versus  $E(P - T)$  (see the text).




**Figure 4.** The expelled masses  $\mathcal{M}(H_0 = 60)$  and  $\mathcal{M}(\text{EPM})$  versus  $\mathcal{M}(P - T)$  (see the text).

distance of 14.53 Mpc is a member of the same group as NGC 3198 and NGC 3319, for which Freedman et al. (2001) give a mean Cepheid distance of 13.5 Mpc,  $\xi$  becomes 1.2. Eventually additional SNe IIP with large distances, where the influence of peculiar motions are negligible, will better determine the scatter of the Hubble diagram and allow a meaningful determination of the actual range of  $\xi$ .

We have considered three sets of the physical supernova parameters  $E$ ,  $\mathcal{M}$ , and  $R$ : (i) for the Hubble distances  $D_H$  with  $H_0 = 60 \text{ km s}^{-1} \text{ Mpc}^{-1}$  (Table 2, column 3); (ii) for the EPM distances  $D_{\text{EPM}}$  (Table 2, column 4); (iii) for the plateau-tail calibrated distances  $D_{P-T}$  (Table 3, column 2).

Although the above parameters derived from the EPM-distances are not presented in Table 2, the corresponding  $E$  and  $\mathcal{M}$ -values can be read out of Figs. 3 and 4 which com-



**Figure 5.** The explosion energy–envelope mass diagram for the case of EPM distances  $D_{\text{EPM}}$  from column 4 of Table 2 (asterisks; SNe 1991al and 1992af being excluded) and for the case of the plateau-tail distances  $D_{\text{P-T}}$  from column 2 of Table 3 (black circles). Some SNe are identified (see text).

pare  $E$  and  $\mathcal{M}$  for sets (i) and (ii) with those for set (iii). For seven SNe  $E$  and  $\mathcal{M}$  are rather insensitive to the adopted distances. However for SNe 1986L and 1990E, labelled in Figs. 3 and 4, the deviations from the (P-T)-values are rather large, especially in case of the envelope mass  $\mathcal{M}$ . These SNe differ from others by having a long plateau of (110 – 120) d in combination with still a substantial expansion velocity of  $4000 \text{ km s}^{-1}$ . As a result, their envelope masses  $\mathcal{M}$ , derived from the distances defined by the  $D_H$  and  $D_{\text{EPM}}$  values, exceed those for other SNe. Such a discrepancy for these two SNe is considerably weakened if  $\xi \approx 2$ . Such a high value of  $\xi$  implies that half of the explosion energy is supplied by a source different from the neutron-proton recombination. This may indicate that for massive SNe the envelope mass  $\mathcal{M}$  (in addition to  $\mathcal{M}_{\text{Ni0}}$ ) is involved in the correlation given by Eq. (8).

The random errors of  $E$  and  $\mathcal{M}$  from our approximate Eqs. (1)–(3) are estimated to be about  $\pm 30\%$ . Observational errors especially in the expansion velocity  $u_{\text{ph}}$  and the plateau duration  $\Delta t$  can modify  $E$  and  $\mathcal{M}$  by another factor of 1.3. Thus it seems reasonable to assume a random uncertainty of a factor of  $\sim 1.5$  for the *individual* values of  $E$  and  $\mathcal{M}$  in Tables 2 and 3. The presupernova radii  $R$  are very sensitive to distance errors (cf. Eq. 5) and may carry random errors of a factor of 2. The radii of SNe with large nickel masses like SN 1991al, 1992af and perhaps 1992am may carry additional systematic errors because Eqs. (1)–(3) do not take into account the radioactive heating in a consistent way.

The expelled masses  $\mathcal{M}$  are plotted against the explosion energies  $E$  in Fig. 5 for two cases, i.e. based on EPM and plateau-tail distances. In case of the  $D_{\text{P-T}}$  distances the mean mass of the eight SNe IIP is  $16 \mathcal{M}_{\odot}$  with an rms deviation of only  $3 \mathcal{M}_{\odot}$ . This narrow mass range is contrasted by a wide range of explosion energies of  $(0.5 - 2.7) \times 10^{51} \text{ erg}$ .


The conclusion that there is no correlation between the expelled mass – which is only  $(1.4 - 2) \mathcal{M}_{\odot}$  smaller than the presupernova mass – and the explosion energy is somewhat weakened by the values of  $\mathcal{M}$  and  $E$  based on the EPM distances suggesting a marginal correlation between  $\mathcal{M}$  and  $E$  which is mainly due to only two SNe: 1986L and 1990E.

One can think of a number of parameters which may explain the wide range of explosion energies. It could be rotation and magnetic fields inherited by the collapsing stellar core. It could be also nonspherical jet-like perturbations of a random nature arising from the macroscopic neutrino-driven advection below the accretion shock. Such perturbations could launch the outgoing blast wave earlier when the recombination nuclear energy stored in a hot neutron-proton gas was not yet as large as it should be in the case of spherical symmetry. If this is correct, one may expect that the asphericity of the explosion *anticorrelates* with the explosion energy.


Recently, a promising project has been started (Van Dyk et al. 1999; Smartt et al. 2001, 2002; and references therein) with the ultimate aim to identify the supernova progenitors (presupernovae) or at least to impose conclusive constraints on their masses by inspecting the pre-discovery field of nearby supernovae. In particular, Smartt et al. derived upper mass limits of  $12 \mathcal{M}_{\odot}$  and  $9 \mathcal{M}_{\odot}$  for the progenitors of the SNe 1999em and 1999gi, assuming distances  $D$  for the host galaxies NGC 1637 and NGC 3184 of 7.5 Mpc and 7.9 Mpc, respectively. Note that these *upper* limits depend on  $D$  and have to be adjusted for other values of  $D$  to  $12 \mathcal{M}_{\odot} (D/7.5 \text{ Mpc})^{0.6}$  for SN 1999em and  $9 \mathcal{M}_{\odot} (D/7.9 \text{ Mpc})^{0.6}$  for SN 1999gi. This follows from the fact that the mass-luminosity relation can be approximated as  $L \sim \mathcal{M}^{3.3}$  in the mass interval  $(10 - 15) \mathcal{M}_{\odot}$ . For SN 1999em at  $D_{\text{P-T}} = 11.08 \text{ Mpc}$  (Table 3) follows  $15.2 \mathcal{M}_{\odot}$  as the upper mass limit for the SN 1999em progenitor. Hence, our result  $\mathcal{M} = 15.0 \mathcal{M}_{\odot}$  (Table 3) does not contradict to the observations *as long as*  $D(1999\text{em}) \gtrsim 10 \text{ Mpc}$ . The situation of SN 1999gi is similar. The upper mass limit for  $D(1999\text{gi}) = 14.53 \text{ Mpc}$  (Table 3) is  $\mathcal{M} < 9 \times (14.53/7.9)^{0.6} = 13.0 \mathcal{M}_{\odot}$ , i.e. not in significant contradiction with the  $\mathcal{M}$ -value of  $14.5 \mathcal{M}_{\odot}$  from Table 3. There is no contradiction either with the upper mass limit of  $15^{+5}_{-3} \mathcal{M}_{\odot}$  for the SN 1999gi progenitor imposed recently by Leonard et al. (2002b).

Equations (1)–(3) by LN85, derived from a grid of 23 SNe IIP models covering a wide parameter space, imply a correlation between the absolute magnitude  $M_V$  (and hence luminosity  $L$  – both measured at the mid-point of the plateau) and the expansion velocity  $u_{\text{ph}}$ . The correlation is shown in Figs. 6 and 7 where 23 grid models are shown by black dots; the straight lines are the least-squares fits. In Fig. 7 are also shown the eight observed SNe IIP from Table 3 marked by open circles, their absolute magnitudes  $M_V$  (Table 3, column 2) being calculated from Eq. 4, where the plateau-tail distances  $D_{\text{P-T}}$  were used from Table 3, column 2. These real SNe follow about the slope of the models, but at a fixed value of  $u_{\text{ph}}$  they are fainter by  $\approx 0.6 \text{ mag}$  on average.

Empirically, Hamuy & Pinto (2002) have also found, using the CMB redshift-based distances, such a correlation. The slopes of their least-squares fits are virtually the same



**Figure 6.** The correlation of the luminosity  $L_{41}$  (in units  $10^{41} \text{ erg s}^{-1}$ ) of the mid-point of the plateau with the expansion velocity  $u_{\text{ph}}$  (in  $1000 \text{ km s}^{-1}$ ) for 23 SN models (black dots).



**Figure 7.** The correlation of the absolute magnitude  $M_V$  of the mid-point of the plateau with the expansion velocity  $u_{\text{ph}}$  for 23 SN models (black dots); open circles relate to 9 real SNe including SN 1987A.

as shown in Figs. 6 and 7. Thus our models confirm their finding.

The main conclusion one can draw from Figs. 6 and 7 is that our three-parametric grid of only 23 SNeIIP properly chosen models is ample enough to reproduce the main features of the real SNe.

## 5 CONCLUSIONS

Model calculation by LN83 and LN85 of SNeIIP, leading to Eqs. (1)–(3), are combined with available EPM distances and velocity distances ( $H_0 = 60$ ) to derive the explosion energy  $E$ , the ejected mass  $\mathcal{M}$ , and the presupernova radius  $R$  of 14 SNeIIP. Only the apparent, absorption-corrected magnitude  $V$  and the expansion velocity  $u_{\text{ph}}$  at the mid-point of

the plateau together with its total duration  $\Delta t$  are needed as additional input parameters. The results are presented in Table 2.

Instead of using EPM or velocity distances it is also possible to use the bolometric fluxes observed during the SNIIP tail phase to determine the Ni mass and hence new, independent distances called here plateau-tail distances  $D_{\text{P-T}}$  (cf. Eq. 11). The  $D_{\text{P-T}}$  distances yield new values of  $E$ ,  $\mathcal{M}$ , and  $R$  given in Table 3 for nine SNe which were observed both during their plateau and tail phases. The values of  $E$  and  $\mathcal{M}$ , based on EPM and P-T distances agree well, with the exception of SNe 1986L and 1990E whose masses  $\mathcal{M}$  coming from P-T distances are by a factor of 2 lower than from EPM distances (see Fig. 4).

The P-T distances are larger than the EPM distances by  $\sim 25\%$  on average. The former suggests a value of  $H_0 = 55 \pm 5$ . The main uncertainty of this result comes from the assumption that  $\xi = 1$ , where  $\xi$  is the ratio between the total explosion energy and the energy liberated by the neutron-proton recombination into  $^{56}\text{Ni}$  (cf. Eq. 10). To reduce the P-T distances to the level of the EPM distances, which correspond to  $H_0 = 70$ , an average value of  $\xi = 1.9$  is required. The consequence that about half of the total energy  $E$  comes from other sources than the neutron-proton recombination into  $^{56}\text{Ni}$  seems rather extreme. In fact it is not supported by two SNeIIP (1987A and 1999gi) with independent distance information, which suggest that  $\xi$  is of order of unity. Moreover, very recently Leonard et al. (2003) have obtained a Cepheid distance of  $11.7 \pm 1$  Mpc to NGC 1637 – the host galaxy of SN 1999em, which is by a factor of 1.4 larger than the EPM distance (Table 2). Our  $D_{\text{P-T}}$  distance of 11.1 Mpc to SN 1999em (Table 3) is in a good agreement with this result. If it happens that the same factor is applicable also to the EPM distances to SNe 1986L and 1990E, there will be no need to resort to large  $\xi$ -values, such as  $\xi \approx 2$  (section 4), to remove the discrepancy between  $D_{\text{P-T}}$  and  $D_{\text{EPM}}$  for these SNe.

In conclusion we emphasize the necessity of constructing a new grid of hydrodynamic SNIIP models based on current evolutionary presupernova models and taking into account  $^{56}\text{Ni}$  as an additional parameter in a consistent way. Such a grid would allow to create more precise analytic approximations for a number of correlations between the physical parameters of SNIIP and their observable properties.

The ‘plateau-tail’ method of distance determination needs, of course, further critical analysis requiring a close collaboration between astronomers observing supernovae and theorists modelling their explosions. If the proposed  $E$ – $\mathcal{M}_{\text{Ni}}$  correlation is confirmed it promises to become a tool to explore the mechanism of SN II with the aid of optical and spectroscopical observations.

## ACKNOWLEDGMENTS

It is a pleasure to express my deep gratitude to the Max-Planck-Institut für Astrophysik for hospitality and financial support. The work was supported also by the Swiss Science National Foundation (Grant 2000/061822) and the Russian Foundation for Basic Research (Project 00-02-17230).

I am grateful to G.A. Tammann for fruitful discussions and a big help and to W. Hillebrandt for useful suggestions. I



thank B. Parodi for acquainting me with the FORTRAN code converting  $v_0$  into  $v_{220}$  and M. Hamuy for sending a copy of his Thesis where a good portion of the observational data used in this work was taken from. The anonymous referee is gratefully acknowledged for the constructive critical comments which helped to improve the paper.

## REFERENCES

- Baron E., Branch D., Hauschildt P. H. et al., 2000, *ApJ*, 545, 444  
 Benetti S., Cappellaro E., Turatto M. et al., 1994, *A&A*, 285, 147  
 Bethe H. A., 1996, *ApJ*, 469, 737  
 Chugai N. N., 1991, *MNRAS*, 250, 513  
 Eastman R. G., Schmidt B. P., Kirshner R. P., 1996, *ApJ*, 466, 911  
 Elmhamdi A., Danziger I. J., Chugai N. et al., 2003, *MNRAS*, 338, 939  
 Freedman W. L., Madore B. F., Gibson B. K. et al., 2001, *ApJ*, 553, 47  
 Grassberg E. K., Nadyozhin D. K., 1976, *Ap&SS*, 44, 409  
 Grassberg E. K., Nadyozhin D. K., 1986, *Astron. Zh.*, 63, 1137 (*Sov. Astron.* 30, 670, 1986)  
 Grassberg E. K., Nadyozhin D. K., 1987, *Astron. Zh.*, 64, 1199 (*Sov. Astron.* 31, 629, 1987)  
 Grassberg E. K., Imshennik V. S., Nadyozhin D. K., 1971, *Ap&SS*, 10, 3  
 Hamuy M., 2001, PhD thesis, Univ. Arizona  
 Hamuy M., Pinto P. A., Maza J. et al., 2001, *ApJ*, 558, 615  
 Hamuy M., Pinto P. A., 2002, *ApJ Lett.*, 566, L63  
 Haynes M. P., van Zee L., Hogg D. E. et al., 1998, *AJ*, 115, 62  
 Imshennik V. S., 2002, *Physics of Atomic Nuclei*, 65, 2077  
 Imshennik V. S., Nadyozhin D. K., 1964, *Astron. Zh.*, 41, 829  
 Imshennik V. S., Nadyozhin D. K., 1989, *Sov. Scientific Rev. (Astrophys. Space Rev.)*, v. 8, Part 1, p. 1–147  
 Kraan-Korteweg R. C., 1986, *A&AS*, 66, 255  
 Leonard D. C. W., Fillipenko A. V., Gates E. L. et al., 2002a, *PASP*, 114, 35  
 Leonard, D. C. W., Fillipenko A. V., Li W. et al., 2002b, *AJ*, 124, 2490  
 Leonard D. C., Kanbur S. M., Ngeow C. C., Tanvir N. R., 2003, accepted for publication in *ApJ*; astro-ph/0305259  
 Li W., Fillipenko A. V., Van Dyk S. D. et al., 2002, *PASP*, 114, 403  
 Litvinova I. Yu., Nadyozhin D. K., 1983, *Ap&SS*, 89, 89 (LN83)  
 Litvinova I. Yu., Nadyozhin D. K., 1985, *Sov. Astron. Lett.*, 11, 145 (LN85)  
 Nadyozhin D. K., 1978, *Ap&SS*, 53, 131  
 Nadyozhin D. K., 1994, *ApJS*, 92, 527  
 Nadyozhin D. K., 2003, in Hillebrandt W., Leibundgut B., eds, *Proc. ESO/MPA/MPE Workshop, From Twilight to High-light: The Physics of Supernovae*. Springer-Verlag, Berlin Heidelberg, p. 241  
 Patat E., Barbon R., Cappellaro E., Turatto, M., 1993, *A&AS*, 98, 443  
 Pennypacker C., Perlmutter S., 1989, *IAU Circ.*, No. 4791  
 Popov D. V., 1993, *ApJ*, 414, 712  
 Rauscher T., Heger A., Hoffman R. D., Woosley S. E., 2002, Preprint UCRL-JC-146742, [http://www-phys.llnl.gov/Rsearch/RRSN/nu\\_csbr/neu\\_rate.html](http://www-phys.llnl.gov/Rsearch/RRSN/nu_csbr/neu_rate.html)  
 Ruiz-Lapuente P., Kidger M., Lopez R., Canal R., 1990, *AJ*, 100, 782  
 Schlegel E. M., 2001, *ApJ*, 556, L25  
 Schmidt B. P., Kirshner R. P., Eastman R. G., 1992, *ApJ*, 395, 366  
 Schmidt B. P., Kirshner R. P., Schild R. et al., 1993, *AJ*, 105, 2236  
 Schmidt B. P., Kirshner R. P., Eastman R. G. et al., 1994a, *ApJ*, 432, 42  
 Schmidt B. P., Kirshner R. P., Eastman R. G. et al., 1994b, *AJ*, 107, 1444  
 Smartt S. J., Gilmore G. F., Trentham N., Tout, C. A., Frayn C. M., 2001, *ApJ*, 556, L29  
 Smartt S. J., Gilmore G. F., Trentham N. et al., 2002, *ApJ*, 565, 1089  
 Thim F., Tammann G. A., Saha A. et al., 2003, *ApJ*, 590, 256  
 Turatto M., Cappellaro E., Benetti S., Danziger I. J., 1993, *MNRAS*, 265, 471  
 Van Dyk S. D., Peng C. Y., Barth A. J. et al., 1999, *PASP*, 111, 313  
 Wood R., Andrews P. J., 1974, *MNRAS*, 167, 13  
 Woosley S. E., Weaver T. A., 1995, *ApJS*, 101, 181

This paper has been produced using the Royal Astronomical Society/Blackwell Science  $\LaTeX$  style file.